

Characteristics of our Varieties

Variety	Resistances				Characteristics						Characteristics / Remarks	
	Maturity	Nematodes	Potato Wart Pathotype	PVY *	PLRV *	Tuber Number	Size of Tubers	Storage	Yield	Flesh Colour / Starch Content		may be harvested from
Table Varieties												
Lord	very early	Ro 1	1	2	2	+	big	very good	very high	yellow	Beginning of August	Fairly firm in cooking (Cooking type AB), salad potato. Combines good taste and very high yield.
Justa	very early	Ro 1	1	3	3	+	big	very good	very high	yellow	End of July	Fairly firm in cooking (Cooking type AB), smooth skin. Very high yield.
Fläming	very early	Ro 1	1	1	1	+	big	very good	very high	light yellow	End of July	Red skin, yellow flesh. Firm in cooking (Cooking type A), excellent salad potato.
Ampera	early	Ro 1	-	2	7	+	small	very good	medium	yellow	Mid- August	A high quality potato, fairly firm in cooking (Cooking type AB). Ideal for packers.
Industry												
Ikone	medium early	Ro 1 + 4	1,2	2	3	+	medium	good	high	18 – 20 %	Mid-October	Suitable for mesh production and as a starch variety.
Kuba	medium early	Ro 1 + 4	1, 2, 6, 8, 18	1	4	+	medium	good	high	18 – 21 %	Mid-August	Suitable for mesh production and as a starch variety. Very little susceptibility to virus-caused diseases, combined with excellent wart resistance. Very high yield at early harvest.
Skonto	medium early	Ro 1	1	3	3	+	medium	good	very high	> 24 %	Mid-September	Extremely high starch in content, already present at an early stage.
Elkana	medium early	Ro 1, 2, 3	1	3	5	-	medium/ big	very good	high	20 – 22 %	Mid-September	Safe yield, even under dry conditions. Excellent variety for long-term storage. Very little susceptibility to damage during harvest.
Starch Varieties												
Novano	medium early	Ro 1, 2, 3, 4 Pa 2, 3	1, 2, 6, 18	2	3	+	medium	good	high	approx. 20 %	End of September	Ideal combination of virus-resistance as well as Nematode and Wart resistance. Ideal for areas with problems in potato health!
Jasia	medium late	Ro 1 + 4	1	1	2	+/-	medium/ big	very good	high	> 21 %	Beginning of October	Abundant and very healthy foliage, even Long vegetation time. A good variety for long-term storage.
Gandawa	medium late	Ro 1	1, 2, 6, 18	1	2	+	medium	fairly good	high	21 %	Mid-October	Gandawa combines a broad potato wart resistance with very good virus resistance. Susceptible to Metribuzin.
Skawa	medium late	Ro 1	1	1	3	++	medium	good	very high	22 – 23 %	Mid-October	Very healthy foliage and tubers. Very high yield potential with high starch content in the tubers.

* Note 1 = very low susceptibility (1 – 10 scale)